

**International Workshop on Waldenstrom's Macroglobulinemia (IWWM-12)
(October 17-19, 2024; Prague, Czech Republic)
Abstract Submission Deadline: Wednesday, July 15, 2024, midnight EST**

Title: Safety and efficacy results of a phase 1 study of the novel BCL2 inhibitor sonrotoclax (BGB-11417) for relapsed/refractory Waldenström macroglobulinemia

Authors: Ramón García Sanz,¹ Chan Cheah,²⁻⁴ Constantine Tam,^{5,5} Mazyar Shadman,⁷ Sophie Leitch,⁸ Christopher D'Angelo,⁹ Lydia Scarfo,¹⁰ Yiqian Fang,¹¹ Sheel Patel,¹² Wei Ding,¹² Haiyi Guo,¹³ Peter Browett,¹⁴

Affiliations: ¹Hospital General Universitario Gregorio Marañón, Madrid, Spain; CIBERONC, Universidad Complutense de Madrid; ²Sir Charles Gairdner Hospital and Pathwest Laboratory Medicine, Nedlands, WA, Australia; ³Medical School, University of Western Australia, Crawley, WA, Australia; ⁴Linear Clinical Research, Nedlands, WA, Australia; ⁵Alfred Hospital, Melbourne, VIC, Australia; ⁶Alfred Hospital, Melbourne, VIC, Australia; ⁷Fred Hutchinson Cancer Research Center, Seattle, WA, USA; ⁸ Te Whatu Ora, Health New Zealand, Waitemata, Auckland, New Zealand; ⁹University Of Nebraska Medical Center, Omaha, NE, USA; ¹⁰Università Vita Salute and IRCCS Ospedale San Raffaele, Milan, Italy; ¹¹BeiGene (Beijing) Co, Ltd, Beijing, China; ¹²BeiGene USA, Inc, San Mateo, CA, USA; ¹³BeiGene (Shanghai) Co, Ltd, Shanghai, China; ¹⁴Auckland City Hospital, Grafton, Auckland, New Zealand

Abstract body: 409/500 words (Graphics are optional and do not count toward limit; no structure requirements)

Introduction: Sonrotoclax (BGB-11417), a novel BCL2 inhibitor, is a more selective and pharmacologically potent inhibitor of BCL2 than venetoclax. Updated safety and

efficacy of sonrotoclax in patients with relapsed/refractory (R/R) Waldenström's macroglobulinemia (WM) from the BGB-11417-101 (NCT04277637) study are presented.

Methods: Patients with R/R WM (≥ 1 prior therapy) received sonrotoclax (planned dose escalation: 80, 160, 320, or 640 mg QD) with dose ramp-up to mitigate tumor lysis syndrome (TLS) risk. The primary endpoint was safety per CTCAE v5.0. The secondary endpoint was ORR (minor response [MR] or better per modified Owens 2013 criteria). TLS was assessed per Howard 2011 criteria.

Results: As of Oct 31, 2023, 17 patients with R/R WM were enrolled (80 mg, n=6; 160 mg, n=8; 320 mg, n=3). Patients had a median age of 68 years (range, 48-87 years) and a median of 2 (range, 2-9) prior treatments. Ten patients had a prior BTK inhibitor (noncovalent, n=1; covalent, n=9) and 14 had prior anti-CD20. Median follow-up was 10.6 (range, 1-24) months. Six patients discontinued treatment (progressive disease [PD], n=4; AEs, n=2 [multifocal neurological syndrome and COVID-19, neither were considered related to sonrotoclax by the investigator]), and 4 patients died on study (PD, n=2; COVID-19 pneumonia, n=1; pneumonia, n=1). TEAEs occurring in $\geq 20\%$ of patients were anemia (n=6; 35%), COVID-19 (n=6; 35%), pyrexia (n=5; 29%), neutropenia (n=4; 24%), and pruritus (n=4; 24%). Anemia was the most common grade ≥ 3 TEAE (n=4; 24%). One patient in the 160-mg dose group experienced a DLT of grade 3 febrile neutropenia, which resolved after 2 days without dose reducing during ramp-up (day 2, 10 mg sonrotoclax). No TLS occurred up to the highest dose tested (320 mg). No atrial or ventricular fibrillation occurred. No deaths or AEs leading to treatment discontinuation were determined by the investigator to be treatment related. In 17 response-evaluable patients, across dose levels, overall, major, and very good partial response (VGPR) rates were 76% (13/17), 41% (7/17), and 12% (2/17), respectively (**Figure**). In 7 patients who had a BTK inhibitor as their last treatment, the ORR was 70% (MR, n=2; PR, n=1; VGPR, n=2). In the 320-mg cohort, the median follow-up was 3.3 (range, 1-5) months and ORR was 100% (n=3), with 1 VGPR.

Conclusions: Sonrotoclax monotherapy was well tolerated and had encouraging preliminary antitumor activity in this heavily pretreated R/R WM population. Based on

these findings, further evaluation of sonrotocax monotherapy in patients with R/R WM is ongoing in a potentially pivotal phase 2 study.

